



# Functions of double-stranded RNA-binding domains in nucleocytoplasmic transport

Silpi Banerjee, Pierre Barraud

## ► To cite this version:

Silpi Banerjee, Pierre Barraud. Functions of double-stranded RNA-binding domains in nucleocytoplasmic transport. *RNA Biology*, 2014, 11 (10), pp.1226 - 1232. 10.4161/15476286.2014.972856 . hal-01110528v2

**HAL Id: hal-01110528**

**<https://hal.science/hal-01110528v2>**

Submitted on 28 May 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Functions of double-stranded RNA-binding domains in nucleocytoplasmic transport

Silpi Banerjee<sup>a</sup> & Pierre Barraud<sup>b,c,d,\*</sup>

<sup>a</sup>Department of Chromosome Biology, Max F. Perutz Laboratories, University of Vienna, A-1030 Vienna, Austria

<sup>b</sup>Centre national de la recherche scientifique, Laboratoire de cristallographie et RMN biologiques, FR-75006 Paris, France

<sup>c</sup>Université Paris Descartes, Sorbonne Paris Cité, Laboratoire de cristallographie et RMN biologiques, FR-75006 Paris, France

<sup>d</sup>Institute of Molecular Biology and Biophysics, Eidgenössische Technische Hochschule Zürich, CH-8093 Zürich, Switzerland

---

## Abstract

The double-stranded RNA-binding domain (dsRBD) is a small protein domain found in eukaryotic, prokaryotic and viral proteins, whose central property is to bind to double-stranded RNA (dsRNA). Aside from this major function, recent examples of dsRBDs involved in the regulation of the sub-cellular localization of proteins, suggest that the participation of dsRBDs in nucleocytoplasmic trafficking is likely to represent a widespread auxiliary function of this type of RNA-binding domain. Overall, dsRBDs from proteins involved in many different biological processes have been reported to be implicated in nuclear import and export, as well as cytoplasmic, nuclear and nucleolar retention. Interestingly, the function of dsRBDs in nucleocytoplasmic trafficking is often regulated by their dsRNA-binding capacity, which can either enhance or impair the transport from one compartment to another. Here, we present and discuss the emerging function of dsRBDs in nucleocytoplasmic transport.

**To cite this article:** S. Banerjee & P. Barraud *RNA Biology* 11 (2014) 1226–1232.

**Full text at the editor website:** <http://www.tandfonline.com/doi/full/10.4161/15476286.2014.972856>

---

## 1. Introduction

RNA performs a multitude of essential functions within the cell, many of which are achieved via interaction with proteins. For instance, many prokaryotic, eukaryotic and viral proteins interact with double-stranded RNA (dsRNA). This type of protein-RNA interaction regulates diverse biological processes such as response to viral infection, gene silencing through RNA interference pathways, RNA processing, regulation of translation, mRNA

editing, RNA export and mRNA localization.<sup>1-4</sup> Proteins interacting with dsRNA often sense the double-helix RNA structure through one or multiple double-stranded RNA-binding domains (dsRBDs). The dsRBD (also referred to as dsRBM for double-stranded RNA-binding motif) is a conserved protein domain of approximately 65-70 amino acids in length, which binds double-stranded or highly structured RNAs. It was first recognized as a conserved protein domain from similarities between *Drosophila* Staufen, human TAR-RNA binding protein (TRBP) and *Xenopus laevis* RNA-binding protein A (XlrpA).<sup>5</sup> The central function of dsRBDs is to bind to dsRNA regions. This is primarily achieved by recognizing the RNA shape, such as for in-

---

\* Corresponding author

Email address: [pierre.barraud@parisdescartes.fr](mailto:pierre.barraud@parisdescartes.fr)  
(Pierre Barraud).

stance the shape of a regular A-form RNA helix or the shape of an RNA hairpin, even though some dsRBDs can bind to dsRNA in a sequence-specific manner (see refs 4, 6-8 and below). Apart from this major function, recent examples of dsRBDs, often with protein-protein interaction properties, have been reported to participate in the regulation of the sub-cellular localization of proteins. This suggests that the participation of dsRBDs in nucleocytoplasmic trafficking is likely to represent a widespread auxiliary function of this type of RNA-binding domain. Among this increasing list of examples, we recently uncovered how the folding of one dsRBD of the human RNA-editing enzyme ADAR1 is assembling a bimodular nuclear localization signal responsible for the active transport of the protein towards the nucleus.<sup>9</sup> In this mini-review, after a brief description of RNA recognition by dsRBDs, we present and discuss the emerging function of dsRBDs in nucleocytoplasmic trafficking in the light of several examples that were recently disclosed.

## 2. Primary function of dsRBDs: dsRNA recognition

Although we will briefly highlight essential features of dsRNA recognition by dsRBDs, interested readers are referred to more comprehensive reviews on these aspects.<sup>1,4,10</sup> dsRBDs can be found in all kingdoms of life and are the second most abundant RNA binding domain,<sup>1</sup> after the RNA recognition motif (RRM), which is a well-characterized single-stranded RNA binding domain.<sup>11</sup> Proteins harboring the dsRBD vary considerably in function, depending on their catalytic domains or interaction partners. So far, the structures of more than 30 dsRBDs have been reported with a conserved  $\alpha 1$ -L1- $\beta 1$ -L2- $\beta 2$ -L3- $\beta 3$ -L4- $\alpha 2$  topology, where L specifies a loop (Figure 1A).<sup>4,12</sup> The second dsRBD of XlrbpA forms the prototype of canonical dsRBDs with the two  $\alpha$ -helices packed against the three-stranded anti-parallel  $\beta$ -sheet.<sup>13,14</sup> In addition, structures of dsRBDs in complex with dsRNA substrates revealed the canonical mode of RNA recognition by dsRBDs.<sup>6,15-17</sup> Molecular recognition is accomplished via three distinct regions of interaction: helix  $\alpha 1$  and the loop between  $\beta 1$  and  $\beta 2$  (L2) contact dsRNA minor grooves at one turn of interval, whereas the N-terminal tip of helix  $\alpha 2$  contacts the dsRNA phosphate backbone across the major groove. Substrate recognition is achieved by virtue of solvent exposed residues in helix  $\alpha 1$ , the GPxH

motif in the  $\beta 1$ - $\beta 2$  loop, and the positively charged KKxAK motif at the N-terminal tip of helix  $\alpha 2$  (Figure 1A). Since the majority of dsRBD-RNA interactions involve direct contacts with the 2'-hydroxyl groups of the ribose sugar rings and direct- or water-mediated contacts with non-bridging oxygen of the phosphodiester backbone, the dsRBD is commonly described as a non-sequence specific RNA binding domain, which perfectly adapts to the shape of the A-form RNA helix.<sup>1,2,18</sup> In addition to this early description, it became clear that some dsRBDs display a pronounced preference for stem-loop structures, with the loop being often a primary determinant of RNA substrate recognition,<sup>8,15,17,19</sup> and that some dsRBDs perform a base-specific readout in the minor-groove via sequence specific contacts involving helix  $\alpha 1$  and the  $\beta 1$ - $\beta 2$  loop.<sup>4,7,20</sup> One should therefore be aware that, beyond the A-form RNA helix, dsRBDs are sensitive to additional RNA features, such as apical loops, base-pair mismatches, bulges and nucleotide sequence.

## 3. dsRBDs and nuclear import

Some of the less conserved dsRBDs (also referred to as type-B dsRBDs)<sup>21</sup> have been shown to be involved in protein localization, while others are required for homo- and/or hetero-dimerization.<sup>12,22,23</sup> An interesting example of a type-B dsRBD involved in localization is the C-terminal dsRBD of human Dicer. Although processing of Dicer's substrates occurs in the cytoplasm of mammalian cells, several reports suggest that Dicer may also have additional roles in the nucleus.<sup>24-27</sup> In support of a nuclear function of Dicer, the C-terminal dsRBD of human Dicer was recently shown to harbor an atypical nuclear localization signal (NLS), composed of a cluster of positively charged residues on the surface of the folded dsRBD (Figure 1B).<sup>28</sup> The dsRBD-dependent import of Dicer follows different pathways and is mediated by Importin- $\beta$ , Importin-7, and Importin-8.<sup>28</sup> However, in the full-length protein, the dsRBD-NLS is masked and Dicer localizes primarily to the cytoplasm. Therefore, even though the dsRBD-NLS mediates nuclear import, human Dicer constantly shuttles between the cytoplasm and the nucleus, and is primarily localized to the cytoplasm under steady-state conditions.<sup>28</sup> The shuttling properties of the dsRBD-NLS might be regulated by another domain of the protein situated in its N-terminal part, namely the helicase domain, and also by the presence of dsRNA.<sup>28</sup> It will there-

fore be interesting to investigate whether the localization of human Dicer is sensitive to variation of external conditions, a layer of regulation that was shown to enable a rapid response to stress conditions in the fission yeast.<sup>29</sup> We recently uncovered the molecular basis for the dsRBD-mediated nuclear import of the RNA-editing enzyme ADAR1.<sup>9</sup> Mammalian ADAR1 is a nucleocytoplasmic shuttling protein containing three dsRBDs. Nuclear import of ADAR1 is mediated by Transportin-1 (Trn1), a member of the karyopherin- $\beta$  family, *via* an atypical NLS that overlaps the third dsRBD of the protein. This atypical NLS shows no similarity to known Trn1 NLSs, commonly referred to as PY-NLS.<sup>30-33</sup> The solution structure of the ADAR1-dsRBD3 revealed an extended dsRBD fold with an additional  $\alpha$ -helix at its N-terminal part (Figure 1C).<sup>9</sup> This N-terminal extension radically changes the relative position of the flexible N- and C-terminal fragments flanking the dsRBD: it brings the N- and C-terminal flanking regions in close proximity (Figure 1C). The two flexible fragments flanking the folded dsRBD constitute two essential modules involved in the interaction with the import receptor Trn1 and thus the nuclear import of ADAR1. The two flanking modules form a so-called bimodular NLS where the two active modules are interrupted by a small folded domain. The intervening dsRBD was shown to only act as a scaffolding domain that properly positions the flanking modules for an effective interaction with Trn1 (Figure 1D). Indeed, functional bimodular NLSs could be designed by replacing the ADAR1-dsRBD3 with another unrelated dsRBD, or even with a small peptide linker that maintained the relative positioning of the N- and C-terminal modules. This clearly indicates that the dsRBD itself is not required for Trn1 interaction, but only helps juxtapose the two NLS-modules that are distantly spaced in the primary structure.<sup>9</sup> In addition, this impressive molecular construction was shown to act as an RNA-sensing NLS that can be switched *on* and *off*, depending on the presence of dsRNA associated with the dsRBD.<sup>9</sup> Indeed, the modulation of the import activity of ADAR1 directly depends on the RNA-binding status of the dsRBD-NLS.<sup>9,33</sup> The dsRBD-NLS presents two functional and non-overlapping interaction surfaces: (i) a functional dsRNA-binding interface that involves the canonical elements of dsRBDs for dsRNA binding; and (ii) a functional Trn1-binding interface that consists of the N- and C-terminal modules flanking the folded dsRBD (Figure 1E). Even though these two

interfaces are distinct, the dsRBD-NLS cannot bind to dsRNA and Trn1 simultaneously, most probably for steric reasons, and therefore constitutes an RNA-sensing NLS.<sup>9</sup> Conversely, RNA-binding enhances nuclear export of ADAR1,<sup>33</sup> which overall positions RNA as a central regulator of ADAR1 shuttling. The fact that dsRNA stimulates nuclear export suggests that ADAR1 might leave the nucleus bound to substrate RNAs, such as precursors of micro-RNAs. The binding of the dsRBD-NLS to Trn1 in the cytoplasm could then help the dissociation of ADAR1-bound RNAs and ensure the reimport of ADAR1 free of RNAs to the nucleus, thereby preventing futile export cycles. In other words, binding of RNA to the RNA-sensing NLS of ADAR1 acts as a control of the directionality of transport, and prevents ADAR1 from carrying RNAs back into the nucleus.

#### 4. dsRBDs and nuclear export

In eukaryotes, the segregation of nuclear and cytoplasmic compartments necessitates the continuous transport of large macromolecules across the nuclear envelop for the proper cellular function.<sup>34,35</sup> Many post-transcriptional functions of RNAs occur in the cytoplasm, far from their nuclear site of biogenesis. Importantly, dsRBDs have been reported to be involved in the export of proteins and/or RNAs through direct or indirect interactions with export factors such as Exportin-5 (Exp5) (see references below). Exp5 is a member of the human karyopherin- $\beta$  family, which shuttles cargo macromolecules from the nucleus to the cytoplasm in a Ran-dependent manner. dsRBD-dependent nuclear export by Exp5 has been reported for many dsRBD-containing proteins such as the elongation factor 1A (eEF1A), the interleukin enhancer binding factor-3 (ILF3), the nuclear factor 90 (NF90), Staufen proteins, and the short isoform of the human editing enzyme ADAR1.<sup>33,36-40</sup> Whether dsRBDs bind to Exp5 directly or *via* a dsRNA bridge seems to vary from one protein to another, and both situations have been reported in the literature (see the three examples below).<sup>33,36-38</sup> For instance, in human ILF3, the second dsRBD of the protein was first proposed to mediate nuclear export through a direct interaction with Exp5,<sup>38</sup> but this interaction, and therefore the nuclear export of the protein, was latter shown to be mediated by the binding of dsRBDs to dsRNA.<sup>36,37</sup> Staufen proteins represent another interesting example of a dsRBD involved in nuclear

export. Staufer localizes mRNAs in *Drosophila*, and is conserved from worms to humans. Staufer proteins possess 4 to 5 dsRBDs depending on the species and protein isoform. It has been reported that nuclear export of mammalian Staufer2 occurs *via* the interaction of the third dsRBD of the protein with Exp5.<sup>39,40</sup> This interaction is abolished in a mutant deficient in RNA-binding, which supports the model of a dsRNA bridge for this dsRBD to interact with Exp5.<sup>39,40</sup> Conversely, the Exp5-mediated nuclear export of the shorter isoform of ADAR1 depends on the interaction of Exp5 with the first dsRBD of the protein, an interaction that is enhanced in the presence of RNA, but persists upon RNase treatment.<sup>33</sup> Therefore, although dsRNA stimulates nuclear export of ADAR1, it seems not essential and binding of dsRBDs to Exp5 seemingly occurs directly.<sup>33</sup> In any case, although the implication of dsRBDs in nuclear export of proteins is often mediated by their RNA-binding properties, their function in nuclear export is central for the subcellular localization of many dsRBD-containing proteins.

## 5. dsRBDs in nuclear or cytoplasmic retention

Another mechanism influencing the subcellular localization of proteins is through modulation of the binding of these proteins to specific nuclear or cytoplasmic factors that anchor, or in other words retain the protein of interest within a specific cytoplasmic or nuclear compartment. The elements involved in the interaction with the anchoring factors are called retention signals, and in the case of dsRBD-containing proteins, dsRBDs have been shown to participate in both nuclear and cytoplasmic retention (see below and references therein). Indeed, similarly to human Dicer, for which the C-terminal dsRBD was shown to be involved in the regulation of the sub-cellular localization of the protein, and especially its transient nuclear import (see ref 28, and text above), the C-terminal dsRBD of the yeast *Schizosaccharomyces pombe* Dicer (Dcr1) is also critical for the nuclear localization of the protein.<sup>41</sup> In contrast to human Dicer, *S. pombe* Dcr1 is predominantly nuclear and localizes at the nuclear periphery, near the nuclear pore complexes (NPCs).<sup>41</sup> Similarly to human Dicer, Dcr1 has the potential to shuttle between the nucleus and cytoplasm, and its C-terminal dsRBD orchestrates its nuclear retention.<sup>41,42</sup> Interestingly, Dcr1's dsRBD folds as an extended dsRBD with an essential C-

terminal extension composed of a short  $\alpha$ -helical turn followed by a zinc-coordination motif of the CHCC type (Figure 1F).<sup>42</sup> Strikingly, only the last two ligands (chCC) are found within the extension, whereas the first two ligands (CHcc) are part of the dsRBD fold itself (respectively in loops L1 and L3). With this particular structure, the extended dsRBD presents a protein-protein interaction surface (Figure 1G), which mediates nuclear retention of Dcr1, by interacting with an unidentified nuclear protein that probably drives its localization at the nuclear periphery.<sup>41,42</sup> Importantly, the extended dsRBD of Dcr1 is a temperature sensitive domain, which acts as a thermoswitch, where elevated temperature above 34-38 °C result in an alteration of the nuclear retention properties of the dsRBD. As a result, Dcr1 localizes in the cytoplasm at elevated temperature, with important consequences for silencing and RNAi-mediated response to stress conditions.<sup>29</sup> Additionally, cytoplasmic retention mediated by dsRBDs has also been reported in a few proteins. For instance, in mammalian Staufer1, dsRBD3 plays critical roles in the cytoplasmic retention of the protein.<sup>43</sup> This retention activity is enhanced in the presence of dsRBD2, and noteworthy, independent of their RNA-binding capacity.<sup>43</sup> Conversely, cytoplasmic retention of the *Xenopus* transcription factor CBTF was also shown to require both its' dsRBDs, but is directly driven by the dsRNA-binding capacity of these domains.<sup>44</sup> Finally, within the nucleus, several dsRBDs have been shown to guide proteins to the nucleolus and to mediate their retention in this compartment. This property was reported for many distinct proteins, such as XlrpA,<sup>45</sup> ADAR proteins,<sup>46-48</sup> and PKR.<sup>49</sup> The targeting of these proteins to the nucleolus is likely driven by their ability to bind to ribosomal RNA.<sup>45-49</sup> For most proteins, it is largely unknown whether the nucleolar localization only reflects their dsRNA-binding capacity, or whether it has a direct biological function.

## 6. Conclusion and perspectives

In this mini-review, we underlined the fact that many dsRBDs play a critical role in the subcellular localization of the protein they belong to. This small and compact domain has been reported to mediate nuclear import and export, as well as cytoplasmic, nuclear, and even nucleolar retention (see Figure 2 for a schematic representation of the diverse functions of dsRBDs in nucleocytoplasmic transport

and for a list of the currently known examples of proteins for which dsRBDs are involved in their sub-cellular localization). Attractively, in the few cases where involvements of dsRBDs in nucleocytoplasmic transport were investigated structurally, extensions to the dsRBD fold have often been described, and these N- or C-terminal extensions were shown to either directly or indirectly participate in the regulation of the protein nucleocytoplasmic distribution.<sup>9,42</sup> In addition, these extended dsRBDs fully retain the capacity to bind to dsRNA, suggesting that extending the dsRBD fold would be a widespread strategy to acquire additional functions without losing the dsRNA-binding activity. More generally, the function of dsRBDs in trafficking is often regulated by their dsRNA-binding capacity, which can either enhance or impair the transport from one compartment to another. For instance, in the case of ADAR1, the competition between the import receptor Trn1 and dsRNA allows the nuclear import to be switched *on* and *off* depending on the presence of RNA associated with the dsRBD-NLS.<sup>9</sup> In our opinion, combining folded functional domains and localization signals is a very attractive molecular construction enabling a myriad of possibilities for the subtle regulation of cellular localization. In a near future, we believe that the list of proteins with a dsRBD involved in subcellular transport and localization will keep growing, and that new aspects of the function of dsRBDs in nucleocytoplasmic transport will surely be uncovered.


## Acknowledgments

We would like to thank Professors Michael Jantsch and Frédéric Allain for helpful discussions and critiques of this manuscript. P.B. is supported by the CNRS and was supported at the ETH by the Postdoctoral ETH Fellowship Program and the Novartis Foundation, formerly the Ciba-Geigy Jubilee Foundation. S.B. was supported by the Austrian Science Foundation SFB4313.

## 7. References

1. Fierro-Monti I, Mathews MB. Proteins binding to duplexed RNA: one motif, multiple functions. *Trends in Biochemical Sciences* 2000; 25:241.
2. Saunders LR, Barber GN. The dsRNA binding protein family: critical roles, diverse cellular functions. *Faseb J* 2003; 17:961-83.
3. Tian B, Bevilacqua PC, Diegelman-Parente A, Mathews MB. The double-stranded-RNA-binding motif: interference and much more. *Nat Rev Mol Cell Biol* 2004; 5:1013-23.
4. Masliah G, Barraud P, Allain FH. RNA recognition by double-stranded RNA binding domains: a matter of shape and sequence. *Cell Mol Life Sci* 2013; 70:1875-95.
5. St Johnston D, Brown NH, Gall JG, Jantsch M. A conserved double-stranded RNA-binding domain. *Proc Natl Acad Sci U S A* 1992; 89:10979-83.
6. Ryter JM, Schultz SC. Molecular basis of double-stranded RNA-protein interactions: structure of a dsRNA-binding domain complexed with dsRNA. *EMBO J* 1998; 17:7505-13.
7. Steff R, Oberstrass FC, Hood JL, Jourdan M, Zimmermann M, Skrisovska L, et al. The solution structure of the ADAR2 dsRBM-RNA complex reveals a sequence-specific readout of the minor groove. *Cell* 2010; 143:225-37.
8. Wang ZH, Hartman E, Roy K, Chanfreau G, Feigon J. Structure of a Yeast RNase III dsRBD Complex with a Noncanonical RNA Substrate Provides New Insights into Binding Specificity of dsRBDs. *Structure* 2011; 19:999-1010.
9. Barraud P, Banerjee S, Mohamed WI, Jantsch MF, Allain FH-T. A bimodular nuclear localization signal assembled via an extended double-stranded RNA-binding domain acts as an RNA-sensing signal for transportin 1. *Proceedings of the National Academy of Sciences* 2014; 111:E1852-E61.
10. Chang KY, Ramos A. The double-stranded RNA-binding motif, a versatile macromolecular docking platform. *The FEBS journal* 2005; 272:2109-17.
11. Maris C, Dominguez C, Allain FH. The RNA recognition motif, a plastic RNA-binding platform to regulate post-transcriptional gene expression. *The FEBS journal* 2005; 272:2118-31.
12. Gleghorn ML, Maquat LE. 'Black sheep' that don't leave the double-stranded RNA-binding domain fold. *Trends Biochem Sci* 2014.
13. Kharrat A, Macias MJ, Gibson TJ, Nilges M, Pastore A. Structure of the dsRNA binding domain of E. coli RNase III. *EMBO J* 1995; 14:3572-84.
14. Bycroft M, Grunert S, Murzin AG, Proctor M, St Johnston D. NMR solution structure of a dsRNA binding domain from Drosophila stau protein reveals homology to the N-terminal domain of ribosomal protein S5. *EMBO J* 1995; 14:3563-71.
15. Ramos A, Grunert S, Adams J, Micklem DR, Proctor MR, Freund S, et al. RNA recognition by a Stau protein double-stranded RNA-binding domain. *EMBO J* 2000; 19:997.
16. Gan J, Tropea JE, Austin BP, Court DL, Waugh DS, Ji X. Structural insight into the mechanism of double-stranded RNA processing by ribonuclease III. *Cell* 2006; 124:355-66.
17. Wu H, Henras A, Chanfreau G, Feigon J. Structural basis for recognition of the AGNN tetraloop RNA fold by the double-stranded RNA-binding domain of Rnt1p RNase III. *Proc Natl Acad Sci U S A* 2004; 101:8307-12.
18. Steff R, Skrisovska L, Allain FH. RNA sequence- and shape-dependent recognition by proteins in the ribonucleoprotein particle. *EMBO reports* 2005; 6:33-8.
19. Liang YH, Lavoie M, Comeau MA, Abou Elela S, Ji X. Structure of a eukaryotic RNase III postcleavage complex reveals a double-ruler mechanism for substrate selection. *Molecular cell* 2014; 54:431-44.


20. Barraud P, Allain FH. ADAR proteins: double-stranded RNA and Z-DNA binding domains. *Current topics in microbiology and immunology* 2012; 353:35-60.
21. Doyle M, Jantsch MF. New and old roles of the double-stranded RNA-binding domain. *Journal of Structural Biology* 2002; 140:147-53.
22. Hitti EG, Sallacz NB, Schoft VK, Jantsch MF. Oligomerization activity of a double-stranded RNA-binding domain. *FEBS Lett* 2004; 574:25-30.
23. Micklem DR, Adams J, Grunert S, St Johnston D. Distinct roles of two conserved Staufen domains in oskar mRNA localization and translation. *The EMBO journal* 2000; 19:1366-77.
24. Sinkkonen L, Hugenschmidt T, Filipowicz W, Svoboda P. Dicer is associated with ribosomal DNA chromatin in mammalian cells. *PLoS one* 2010; 5:e12175.
25. Gullerova M, Proudfoot NJ. Convergent transcription induces transcriptional gene silencing in fission yeast and mammalian cells. *Nat Struct Mol Biol* 2012; 19:1193-201.
26. White E, Schlackow M, Kamieniarz-Gdula K, Proudfoot NJ, Gullerova M. Human nuclear Dicer restricts the deleterious accumulation of endogenous double-stranded RNA. *Nat Struct Mol Biol* 2014; 21:552-9.
27. Gagnon KT, Li L, Chu Y, Janowski BA, Corey DR. RNAi factors are present and active in human cell nuclei. *Cell reports* 2014; 6:211-21.
28. Doyle M, Badertscher L, Jaskiewicz L, Guttinger S, Jurado S, Hugenschmidt T, et al. The double-stranded RNA binding domain of human Dicer functions as a nuclear localization signal. *RNA* 2013; 19:1238-52.
29. Woolcock KJ, Stunnenberg R, Gaidatzis D, Hotz HR, Emmerth S, Barraud P, et al. RNAi keeps Atf1-bound stress response genes in check at nuclear pores. *Genes Dev* 2012; 26:683-92.
30. Eckmann CR, Neunteufl A, Pfaffstetter L, Jantsch MF. The human but not the *Xenopus* RNA-editing enzyme ADAR1 has an atypical nuclear localization signal and displays the characteristics of a shuttling protein. *Mol Biol Cell* 2001; 12:1911-24.
31. Poulsen H, Nilsson J, Damgaard CK, Egebjerg J, Kjems J. CRM1 Mediates the Export of ADAR1 through a Nuclear Export Signal within the Z-DNA Binding Domain. *Molecular and cellular biology* 2001; 21:7862-71.
32. Strehblow A, Hallegger M, Jantsch MF. Nucleocytoplasmic distribution of human RNA-editing enzyme ADAR1 is modulated by double-stranded RNA-binding domains, a leucine-rich export signal, and a putative dimerization domain. *Mol Biol Cell* 2002; 13:3822-35.
33. Fritz J, Strehblow A, Taschner A, Schopoff S, Pasierbek P, Jantsch MF. RNA-regulated interaction of transportin-1 and exportin-5 with the double-stranded RNA-binding domain regulates nucleocytoplasmic shuttling of ADAR1. *Mol Cell Biol* 2009; 29:1487-97.
34. Fried H, Kutay U. Nucleocytoplasmic transport: taking an inventory. *Cell Mol Life Sci* 2003; 60:1659-88.
35. Cook A, Bono F, Jinek M, Conti E. Structural biology of nucleocytoplasmic transport. *Annual review of biochemistry* 2007; 76:647-71.
36. Calado A, Treichel N, Muller EC, Otto A, Kutay U. Exportin-5-mediated nuclear export of eukaryotic elongation factor 1A and tRNA. *The EMBO journal* 2002; 21:6216-24.
37. Gwizdek C, Ossareh-Nazari B, Brownawell AM, Evers S, Macara IG, Dargemont C. Minihelix-containing RNAs mediate exportin-5-dependent nuclear export of the double-stranded RNA-binding protein ILF3. *J Biol Chem* 2004; 279:884-91.
38. Brownawell AM, Macara IG. Exportin-5, a novel karyopherin, mediates nuclear export of double-stranded RNA binding proteins. *J Cell Biol* 2002; 156:53-64.
39. Miki T, Takano K, Yoneda Y. The Role of Mammalian Staufen on mRNA Traffic: a View from Its Nucleocytoplasmic Shuttling Function. *Cell Structure and Function* 2005; 30:51-6.
40. Macchi P, Brownawell AM, Grunewald B, DesGroseillers L, Macara IG, Kiebler MA. The brain-specific double-stranded RNA-binding protein Staufen2: nucleolar accumulation and isoform-specific exportin-5-dependent export. *J Biol Chem* 2004; 279:31440-4.
41. Emmerth S, Schober H, Gaidatzis D, Roloff T, Jacobeit K, Buhler M. Nuclear retention of fission yeast dicer is a prerequisite for RNAi-mediated heterochromatin assembly. *Developmental cell* 2010; 18:102-13.
42. Barraud P, Emmerth S, Shimada Y, Hotz H-R, Allain FH-T, Bühler M. An extended dsRBD with a novel zinc-binding motif mediates nuclear retention of fission yeast Dicer. *EMBO J* 2011; 30:4223-35.
43. Martel C, Macchi P, Furic L, Kiebler MA, DesGroseillers L. Staufen1 is imported into the nucleolus via a bipartite nuclear localization signal and several modulatory determinants. *Biochem J* 2006; 393:245-54.
44. Brzostowski J, Robinson C, Orford R, Elgar S, Scarlett G, Peterkin T, et al. RNA-dependent cytoplasmic anchoring of a transcription factor subunit during *Xenopus* development. *EMBO J* 2000; 19:3683-93.
45. Eckmann CR, Jantsch MF. Xlrpba, a double-stranded RNA-binding protein associated with ribosomes and heterogeneous nuclear RNPs. *J Cell Biol* 1997; 138:239-53.
46. Desterro JM, Keegan LP, Lafarga M, Berciano MT, O'Connell M, Carmo-Fonseca M. Dynamic association of RNA-editing enzymes with the nucleolus. *J Cell Sci* 2003; 116:1805-18.
47. Sansam CL, Wells KS, Emeson RB. Modulation of RNA editing by functional nucleolar sequestration of ADAR2. *Proc Natl Acad Sci U S A* 2003; 100:14018-23.
48. Xu M, Wells KS, Emeson RB. Substrate-dependent contribution of double-stranded RNA-binding motifs to ADAR2 function. *Mol Biol Cell* 2006; 17:3211-20.
49. Tian B, Mathews MB. Functional Characterization of and Cooperation between the Double-stranded RNA-binding Motifs of the Protein Kinase PKR. *Journal of Biological Chemistry* 2001; 276:9936-44.



**Figure 1. Structural aspects of the functions of double-stranded RNA-binding domains.**

(A) Primary function of dsRBDs as dsRNA binders. Cartoon representation of the canonical mode of interaction of dsRBDs with dsRNA (ADAR2-dsRBD1 is shown as an example; PDB code 2l3c).<sup>7</sup>  $\beta$ -strands are in yellow and  $\alpha$ -helices in blue. The three regions of interaction with dsRNA are in red, with the three conserved lysines of the so-called KKxAK motif shown as sticks. Helix  $\alpha$ 1 and the loop L2 interact with dsRNA minor grooves at one turn of interval. The N-terminal tip of helix  $\alpha$ 2 interacts with the intervening major groove. (B) Cartoon representation of the C-terminal dsRBD of human Dicer (PDB code 3c4b). The atypical dsRBD-NLS of human Dicer is composed of a cluster of positively charged residues represented as sticks in red (namely K1891, R1910 and R1907).<sup>28</sup> Noteworthy, these residues do not belong to the canonical region of interaction with dsRNA. (C) Cartoon representation of the dsRBD-NLS of human ADAR1 (PDB code 2mdr).<sup>9</sup> The atypical dsRBD-NLS of human ADAR1 is composed of the N- and C-terminal modules (in blue and in red, respectively) flanking the folded dsRBD. The additional N-terminal helix (helix  $\alpha$ N, in purple) brings the two modules in close proximity. (D) Schematic representation of the interaction of the ADAR1 dsRBD-NLS with Trn1. The dsRBD core domain is in gray, the N- and C-terminal modules are in blue and in red, respectively, and the additional helix  $\alpha$ N is in purple. Transportin-1 is represented as a gray arch. (E) Schematic representation of the two functional binding interfaces of the ADAR1 dsRBD-NLS. These two binding sites cannot be occupied simultaneously, most probably for steric reasons, and therefore, the dsRBD-NLS constitutes an RNA-sensing NLS.<sup>9</sup> (F) Cartoon representation of the C-terminal dsRBD of *S. pombe* Dicer (Dcr1, PDB code 2l6m).<sup>42</sup>  $\beta$ -strands are in yellow and  $\alpha$ -helices in blue. Dcr1's dsRBD folds as an extended dsRBD with a C-terminal extension composed of a short  $\alpha$ -helical turn (helix  $\alpha$ 3) and a zinc-coordination motif of the CHCC type. The zinc ion is shown as a gray sphere, and the CHCC ligands as blue sticks. (G) The C-terminal extension of Dcr1's dsRBD provides an additional surface, and the residues highlighted in red (namely N1344, Y1348 and S1349) form a protein-protein interaction surface that is required to retain Dcr1 in the nucleus.<sup>42</sup>





**Figure 2. Functions of double-stranded RNA-binding domains in nucleocytoplasmic transport.** dsRBDs (red spheres) are involved in many aspects governing the subcellular localization of proteins. This includes nuclear import and export (symbolized with red arrows), as well as cytoplasmic, nuclear, and nucleolar retention *via* the interaction with anchoring factors (blue polygons). Other types of protein domains are symbolized as spheres of various colors. For each function of dsRBDs in nucleocytoplasmic transport (text in red), the protein examples such as presented in this mini-review are listed below (text in black). This list does not aim at being exhaustive and is likely to increase in the near future.